




World Health Organization


Mortality Country Fact Sheet 2006

Swaziland




Summary	Year	Males	Females	Both sexes
Population (millions)	2005	.5	.5	1.0
Life expectancy (years)	2004	36	39	37
Under-5 mortality (per 1 000 live births)	2004	163	150	156
Adult mortality (per 1 000)	2004	823	741	
Maternal mortality (per 100 000 live births)	2000		370	


Source: World Health Statistics 2006



Under-5 mortality



Source: i) WHO mortality database ii) World Health Statistics 2006



Source: World Health Report 2005

Under-5 mortality: for highest and lowest quintiles
Swaziland, DHS

Category	Lowest	Highest	Ratio
Wealth/assets quintiles
Sex	Males	Females	Ratio

Urban/Rural	Rural	Urban	Ratio

Mother's education quintiles	None	Higher	Ratio

Note: rate per 1 000 live births for 10-year period preceding the survey
Source: DHS



Note: rate for 10-year period preceding the survey
Source: DHS



World Health Organization

Mortality Country Fact Sheet 2006

Causes of death in children under-5

Distribution of causes of death among children under 5 years of age Swaziland, 2000-2003			Annual estimated proportions of death by cause for neonates Swaziland, 2000		
Causes	Deaths ^b (%)	Regional average (%)	Causes	Deaths ^c (%)	Regional average ^c (%)
Total neonatal deaths	100	100	Total neonatal deaths	100	100
Neonatal causes ^a	27	26	Neonatal tetanus	2	9
HIV/AIDS	47	7	Severe infection ^a	25	27
Diarrhoeal diseases	10	17	Birth asphyxia	26	24
Measles	0	4	Diarrhoeal diseases	3	3
Malaria	0	17	Congenital anomalies	8	6
Pneumonia	12	21	Preterm birth ^b	29	23
Injuries	4	2	Others	7	7
Others	1	6			

a. Includes diarrhoea during neonatal period
b. Sum of individual proportions may not add up to 100% due to rounding.

a. Includes deaths from pneumonia, meningitis, sepsis/septicaemia and other infections during the neonatal period.
b. Includes only deaths directly attributed to prematurity and to specific complications of preterm birth such as surfactant deficiency, but not all deaths in preterm infants.
c. Sum of individual proportions may not equal 100% due to rounding.

Causes of Death

Top ten causes of death, all ages Swaziland, 2002			
Causes	Deaths		Years of Life Lost (%)
	(000)	(%)	
All causes	26	100	100
HIV/AIDS	16	64	68
Lower respiratory infections	1	5	6
Tuberculosis	1	4	4
Diarrhoeal diseases	0	2	3
Ischaemic heart disease	0	2	1
Cerebrovascular disease	0	2	1
Perinatal conditions	0	2	2
Protein-energy malnutrition	0	1	2
Road traffic accidents	0	1	1
Chronic obstructive pulmonary disease	0	1	0

Source: [Death and DALY estimates by cause, 2002](http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls)
<http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls>

Life expectancy at birth among males (years) Swaziland, 2004

Life expectancy at birth among females (years) Swaziland, 2004

Maternal mortality ratio (per 100 000 live births) Swaziland, 2000

Legend:
 WHO African Region
 Swaziland

Source: World Health Statistics 2006